

Soy Alert -- Tragedy and Hype

The Third International Soy Symposium

by Sally Fallon & Mary Enig, PhD

First published in *Nexus Magazine*, Volume 7, Number 3, April-May, 2000
© 2000 by Sally Fallon and Mary G. Enig, PhD. All rights reserved.

1. Marketing the Perfect Food
2. Cinderella's Dark Side
3. Soy Protein Isolate
4. The FDA Health Claim
5. Soy and Cancer
6. Phytoestrogens -- Panacea or Poison?
7. Birth Control Pills for Babies
8. Dissension in the Ranks
9. GRAS Status
10. The Next Asbestos?
11. References

Far from being the perfect food, modern soy products contain antinutrients and toxins and they interfere with the absorption of vitamins and minerals.

"Each year, research on the health effects of soy and soybean components seems to increase exponentially. Furthermore, research is not just expanding in the primary areas under investigation, such as cancer, heart disease and osteoporosis; new findings suggest that soy has potential benefits that may be more extensive than previously thought." So writes Mark Messina, PhD, General Chairperson of the Third International Soy Symposium, held in Washington, DC, in November 1999.^[1]

For four days, well-funded scientists gathered in Washington made presentations to an admiring press and to their sponsors -- United Soybean Board, American Soybean Association, Monsanto, Protein Technologies International, Central Soya, Cargill Foods, Personal Products Company, SoyLife, Whitehall-Robins Healthcare and the soybean councils of Illinois, Indiana, Kentucky, Michigan, Minnesota, Nebraska, Ohio and South Dakota.

The symposium marked the apogee of a decade-long marketing campaign to gain consumer acceptance of tofu, soy milk, soy ice cream, soy cheese, soy sausage and soy derivatives, particularly soy isoflavones like genistein and diadzen, the oestrogen-like compounds found in soybeans. It coincided with a US Food and Drug Administration (FDA) decision, announced on October 25, 1999, to allow a health claim for products "low in saturated fat and cholesterol" that contain 6.25 grams of soy protein per serving. Breakfast cereals, baked goods, convenience food, smoothie mixes and meat substitutes could now be sold with labels touting benefits to cardiovascular health, as long as these products contained one heaping teaspoon of soy protein per 100-gram serving.

Marketing the Perfect Food

"Just imagine you could grow the perfect food. This food not only would provide affordable nutrition, but also would be delicious and easy to prepare in a variety of ways. It would be a healthful food, with no saturated fat. In fact, you would be growing a virtual fountain of youth on your back forty." The author is Dean Houghton, writing for *The Furrow*,^[2] a magazine published in 12 languages by John Deere. "This ideal food would help prevent, and perhaps reverse, some of the world's most dreaded diseases. You could grow this miracle crop in a variety of soils and climates. Its cultivation would build up, not deplete, the land . . . this miracle food already exists . . . It's called soy."

Just imagine. Farmers have been imagining -- and planting more soy. What was once a minor crop, listed in the 1913 US Department of Agriculture (USDA) handbook not as a food but as an industrial product, now covers 72 million acres of American farmland. Much of this harvest will be used to feed chickens, turkeys, pigs, cows and salmon. Another large fraction will be squeezed to produce oil for margarine, shortenings and salad dressings.

Advances in technology make it possible to produce isolated soy protein from what was once considered a waste product -- the defatted, high-protein soy chips -- and then transform something that looks and smells terrible into products that can be consumed by human beings. Flavourings, preservatives, sweeteners, emulsifiers and synthetic nutrients have turned soy protein isolate, the food processors' ugly duckling, into a New Age Cinderella.

Lately, this new fairy-tale food has been marketed not so much for her beauty but for her virtues. Early on, products based on soy protein isolate were sold as extenders and meat substitutes -- a strategy that failed to produce the requisite consumer demand. The industry changed its approach. "The quickest way to gain product acceptability in the less affluent society," said an industry spokesman, "is to have the product consumed on its own merit in a more affluent society."^[3] So soy is now sold to the upscale consumer, not as a cheap, poverty food but as a miracle substance that will prevent heart disease and cancer, whisk away hot flushes, build strong bones and keep us forever young. The competition -- meat, milk, cheese, butter and eggs - has been duly demonised by the appropriate government bodies. Soy serves as meat and milk for a new generation of virtuous vegetarians.

Marketing costs money, especially when it needs to be bolstered with "research", but there's plenty of funds available. All soybean producers pay a mandatory assessment of one-half to

one per cent of the net market price of soybeans. The total -- something like US\$80 million annually^[4] -- supports United Soybean's program to "strengthen the position of soybeans in the marketplace and maintain and expand domestic and foreign markets for uses for soybeans and soybean products". State soybean councils from Maryland, Nebraska, Delaware, Arkansas, Virginia, North Dakota and Michigan provide another \$2.5 million for "research".^[5] Private companies like Archer Daniels Midland also contribute their share. ADM spent \$4.7 million for advertising on *Meet the Press* and \$4.3 million on *Face the Nation* during the course of a year.^[6] Public relations firms help convert research projects into newspaper articles and advertising copy, and law firms lobby for favourable government regulations. IMF money funds soy processing plants in foreign countries, and free trade policies keep soybean abundance flowing to overseas destinations.

The push for more soy has been relentless and global in its reach. Soy protein is now found in most supermarket breads. It is being used to transform "the humble tortilla, Mexico's corn-based staple food, into a protein-fortified 'super-tortilla' that would give a nutritional boost to the nearly 20 million Mexicans who live in extreme poverty".^[7] Advertising for a new soy-enriched loaf from Allied Bakeries in Britain targets menopausal women seeking relief from hot flashes. Sales are running at a quarter of a million loaves per week.^[8]

The soy industry hired Norman Robert Associates, a public relations firm, to "get more soy products onto school menus".^[9] The USDA responded with a proposal to scrap the 30 per cent limit for soy in school lunches. The NuMenu program would allow unlimited use of soy in student meals. With soy added to hamburgers, tacos and lasagna, dieticians can get the total fat content below 30 per cent of calories, thereby conforming to government dictates. "With the soy-enhanced food items, students are receiving better servings of nutrients and less cholesterol and fat."

Soy milk has posted the biggest gains, soaring from \$2 million in 1980 to \$300 million in the US last year.^[10] Recent advances in processing have transformed the grey, thin, bitter, beany-tasting Asian beverage into a product that Western consumers will accept -- one that tastes like a milkshake, but without the guilt.

Processing miracles, good packaging, massive advertising and a marketing strategy that stresses the products' possible health benefits account for increasing sales to all age groups. For example, reports that soy helps prevent prostate cancer have made soy milk acceptable to middle-aged men. "You don't have to twist the arm of a 55- to 60-year-old guy to get him to try soy milk," says Mark Messina. Michael Milken, former junk bond financier, has helped the industry shed its hippie image with well-publicised efforts to consume 40 grams of soy protein daily. Now it's OK for stockbrokers to eat soy.

America today, tomorrow the world. Soy milk sales are rising in Canada, even though soy milk there costs twice as much as cow's milk. Soybean milk processing plants are sprouting up in places like Kenya.^[11] Even China, where soy really is a poverty food and whose people want more meat, not tofu, has opted to build Western-style soy factories rather than develop western grasslands for grazing animals.^[12]

Cinderella's Dark Side

The propaganda that has created the soy sales miracle is all the more remarkable because, only a few decades ago, the soybean was considered unfit to eat -- even in Asia. During the Chou Dynasty (1134&endash;246 BC) the soybean was designated one of the five sacred grains, along with barley, wheat, millet and rice. However, the pictograph for the soybean, which dates from earlier times, indicates that it was not first used as a food; for whereas the pictographs for the other four grains show the seed and stem structure of the plant, the pictograph for the soybean emphasises the root structure. Agricultural literature of the period speaks frequently of the soybean and its use in crop rotation. Apparently the soy plant was initially used as a method of fixing nitrogen.^[13]

The soybean did not serve as a food until the discovery of fermentation techniques, some time during the Chou Dynasty. The first soy foods were fermented products like tempeh, natto, miso and soy sauce. At a later date, possibly in the 2nd century BC, Chinese scientists discovered that a purée of cooked soybeans could be precipitated with calcium sulphate or magnesium sulphate (plaster of Paris or Epsom salts) to make a smooth, pale curd -- tofu or bean curd. The use of fermented and precipitated soy products soon spread to other parts of the Orient, notably Japan and Indonesia.

The Chinese did not eat unfermented soybeans as they did other legumes such as lentils because the soybean contains large quantities of natural toxins or "antinutrients". First among them are potent enzyme inhibitors that block the action of trypsin and other enzymes needed for protein digestion. These inhibitors are large, tightly folded proteins that are not completely deactivated during ordinary cooking. They can produce serious gastric distress, reduced protein digestion and chronic deficiencies in amino acid uptake. In test animals, diets high in trypsin inhibitors cause enlargement and pathological conditions of the pancreas, including cancer.^[14]

Soybeans also contain haemagglutinin, a clot-promoting substance that causes red blood cells to clump together.

Trypsin inhibitors and haemagglutinin are growth inhibitors. Weaning rats fed soy containing these antinutrients fail to grow normally. Growth-depressant compounds are deactivated during the process of fermentation, so once the Chinese discovered how to ferment the soybean, they began to incorporate soy foods into their diets. In precipitated products, enzyme inhibitors concentrate in the soaking liquid rather than in the curd. Thus, in tofu and bean curd, growth depressants are reduced in quantity but not completely eliminated.

Soy also contains goitrogens -- substances that depress thyroid function.

Soybeans are high in phytic acid, present in the bran or hulls of all seeds. It's a substance that can block the uptake of essential minerals -- calcium, magnesium, copper, iron and especially zinc -- in the intestinal tract. Although not a household word, phytic acid has been extensively studied; there are literally hundreds of articles on the effects of phytic acid in the current scientific literature. Scientists are in general agreement that grain- and legume-based diets high in phytates contribute to widespread mineral deficiencies in third world

countries.^[15] Analysis shows that calcium, magnesium, iron and zinc are present in the plant foods eaten in these areas, but the high phytate content of soy- and grain-based diets prevents their absorption.

The soybean has one of the highest phytate levels of any grain or legume that has been studied,^[16] and the phytates in soy are highly resistant to normal phytate-reducing techniques such as long, slow cooking.^[17] Only a long period of fermentation will significantly reduce the phytate content of soybeans. When precipitated soy products like tofu are consumed with meat, the mineral-blocking effects of the phytates are reduced.^[18] The Japanese traditionally eat a small amount of tofu or miso as part of a mineral-rich fish broth, followed by a serving of meat or fish.

Vegetarians who consume tofu and bean curd as a substitute for meat and dairy products risk severe mineral deficiencies. The results of calcium, magnesium and iron deficiency are well known; those of zinc are less so.

Zinc is called the intelligence mineral because it is needed for optimal development and functioning of the brain and nervous system. It plays a role in protein synthesis and collagen formation; it is involved in the blood-sugar control mechanism and thus protects against diabetes; it is needed for a healthy reproductive system. Zinc is a key component in numerous vital enzymes and plays a role in the immune system. Phytates found in soy products interfere with zinc absorption more completely than with other minerals.^[19] Zinc deficiency can cause a "spacey" feeling that some vegetarians may mistake for the "high" of spiritual enlightenment.

Milk drinking is given as the reason why second-generation Japanese in America grow taller than their native ancestors. Some investigators postulate that the reduced phytate content of the American diet -- whatever may be its other deficiencies -- is the true explanation, pointing out that both Asian and Western children who do not get enough meat and fish products to counteract the effects of a high phytate diet, frequently suffer rickets, stunting and other growth problems.^[20]

Soy Protein Isolate

Soy processors have worked hard to get these antinutrients out of the finished product, particularly soy protein isolate (SPI) which is the key ingredient in most soy foods that imitate meat and dairy products, including baby formulas and some brands of soy milk.

SPI is not something you can make in your own kitchen. Production takes place in industrial factories where a slurry of soy beans is first mixed with an alkaline solution to remove fibre, then precipitated and separated using an acid wash and, finally, neutralised in an alkaline solution. Acid washing in aluminium tanks leaches high levels of aluminium into the final product. The resultant curds are spray-dried at high temperatures to produce a high-protein powder. A final indignity to the original soybean is high-temperature, high-pressure extrusion processing of soy protein isolate to produce textured vegetable protein (TVP).

Much of the trypsin inhibitor content can be removed through high-temperature processing, but not all. Trypsin inhibitor content of soy protein isolate can vary as much as fivefold.^[21] (In rats, even low-level trypsin inhibitor SPI feeding results in reduced weight gain compared to controls.^[22]) But high-temperature processing has the unfortunate side-effect of so denaturing the other proteins in soy that they are rendered largely ineffective.^[23] That's why animals on soy feed need lysine supplements for normal growth.

Nitrites, which are potent carcinogens, are formed during spray-drying, and a toxin called lysinoalanine is formed during alkaline processing.^[24] Numerous artificial flavourings, particularly MSG, are added to soy protein isolate and textured vegetable protein products to mask their strong "beany" taste and to impart the flavour of meat.^[25]

In feeding experiments, the use of SPI increased requirements for vitamins E, K, D and B12 and created deficiency symptoms of calcium, magnesium, manganese, molybdenum, copper, iron and zinc.^[26] Phytic acid remaining in these soy products greatly inhibits zinc and iron absorption; test animals fed SPI develop enlarged organs, particularly the pancreas and thyroid gland, and increased deposition of fatty acids in the liver.^[27]

Yet soy protein isolate and textured vegetable protein are used extensively in school lunch programs, commercial baked goods, diet beverages and fast food products. They are heavily promoted in third world countries and form the basis of many food giveaway programs.

In spite of poor results in animal feeding trials, the soy industry has sponsored a number of studies designed to show that soy protein products can be used in human diets as a replacement for traditional foods. An example is "Nutritional Quality of Soy Bean Protein Isolates: Studies in Children of Preschool Age", sponsored by the Ralston Purina Company.^[28] A group of Central American children suffering from malnutrition was first stabilised and brought into better health by feeding them native foods, including meat and dairy products. Then, for a two-week period, these traditional foods were replaced by a drink made of soy protein isolate and sugar. All nitrogen taken in and all nitrogen excreted was measured in truly Orwellian fashion: the children were weighed naked every morning, and all excrement and vomit gathered up for analysis. The researchers found that the children retained nitrogen and that their growth was "adequate", so the experiment was declared a success.

Whether the children were actually healthy on such a diet, or could remain so over a long period, is another matter. The researchers noted that the children vomited "occasionally", usually after finishing a meal; that over half suffered from periods of moderate diarrhoea; that some had upper respiratory infections; and that others suffered from rash and fever.

It should be noted that the researchers did not *dare* to use soy products to help the children recover from malnutrition, and were obliged to supplement the soy-sugar mixture with nutrients largely absent in soy products -- notably, vitamins A, D and B12, iron, iodine and zinc.

The FDA Health Claim

The best marketing strategy for a product that is inherently unhealthy is, of course, a health claim.

"The road to FDA approval," writes a soy apologist, "was long and demanding, consisting of a detailed review of human clinical data collected from more than 40 scientific studies conducted over the last 20 years. Soy protein was found to be one of the rare foods that had sufficient scientific evidence not only to qualify for an FDA health claim proposal but to ultimately pass the rigorous approval process."^[29]

The "long and demanding" road to FDA approval actually took a few unexpected turns. The original petition, submitted by Protein Technology International, requested a health claim for isoflavones, the oestrogen-like compounds found plentifully in soybeans, based on assertions that "only soy protein that has been processed in a manner in which isoflavones are retained will result in cholesterol lowering". In 1998, the FDA made the unprecedented move of rewriting PTI's petition, removing any reference to the phyto-oestrogens and substituting a claim for soy protein -- a move that was in direct contradiction to the agency's regulations. The FDA is authorised to make rulings only on substances presented by petition.

The abrupt change in direction was no doubt due to the fact that a number of researchers, including scientists employed by the US Government, submitted documents indicating that isoflavones are toxic.

The FDA had also received, early in 1998, the final British Government report on phytoestrogens, which failed to find much evidence of benefit and warned against potential adverse effects.^[30]

Even with the change to soy protein isolate, FDA bureaucrats engaged in the "rigorous approval process" were forced to deal nimbly with concerns about mineral blocking effects, enzyme inhibitors, goitrogenicity, endocrine disruption, reproductive problems and increased allergic reactions from consumption of soy products.^[31]

One of the strongest letters of protest came from Dr Dan Sheehan and Dr Daniel Doerge, government researchers at the National Center for Toxicological Research.^[32] Their pleas for warning labels were dismissed as unwarranted.

"Sufficient scientific evidence" of soy's cholesterol-lowering properties is drawn largely from a 1995 meta-analysis by Dr James Anderson, sponsored by Protein Technologies International and published in the *New England Journal of Medicine*.^[33]

A meta-analysis is a review and summary of the results of many clinical studies on the same subject. Use of meta-analyses to draw general conclusions has come under sharp criticism by members of the scientific community. "Researchers substituting meta-analysis for more rigorous trials risk making faulty assumptions and indulging in creative accounting," says Sir John Scott, President of the Royal Society of New Zealand. "Like is not being lumped with like. Little lumps and big lumps of data are being gathered together by various groups."^[34]

There is the added temptation for researchers, particularly researchers funded by a company like Protein Technologies International, to leave out studies that would prevent the desired conclusions. Dr Anderson discarded eight studies for various reasons, leaving a remainder of twenty-nine. The published report suggested that individuals with cholesterol levels over 250 mg/dl would experience a "significant" reduction of 7 to 20 per cent in levels of serum cholesterol if they substituted soy protein for animal protein. Cholesterol reduction was insignificant for individuals whose cholesterol was lower than 250 mg/dl.

In other words, for most of us, giving up steak and eating veggieburgers instead will not bring down blood cholesterol levels. The health claim that the FDA approved "after detailed review of human clinical data" fails to inform the consumer about these important details.

Research that ties soy to positive effects on cholesterol levels is "incredibly immature", said Ronald M. Krauss, MD, head of the Molecular Medical Research Program and Lawrence Berkeley National Laboratory.^[35] He might have added that studies in which cholesterol levels were lowered through either diet or drugs have consistently resulted in a greater number of deaths in the treatment groups than in controls -- deaths from stroke, cancer, intestinal disorders, accident and suicide.^[36] Cholesterol-lowering measures in the US have fuelled a \$60 billion per year cholesterol-lowering industry, but have not saved us from the ravages of heart disease.

Soy and Cancer

The new FDA ruling does not allow any claims about cancer prevention on food packages, but that has not restrained the industry and its marketeers from making them in their promotional literature.

"In addition to protecting the heart," says a vitamin company brochure, "soy has demonstrated powerful anticancer benefits . . . the Japanese, who eat 30 times as much soy as North Americans, have a lower incidence of cancers of the breast, uterus and prostate."^[37]

Indeed they do. But the Japanese, and Asians in general, have much higher rates of other types of cancer, particularly cancer of the oesophagus, stomach, pancreas and liver.^[38] Asians throughout the world also have high rates of thyroid cancer.^[39] The logic that links low rates of reproductive cancers to soy consumption requires attribution of high rates of thyroid and digestive cancers to the same foods, particularly as soy causes these types of cancers in laboratory rats.

Just how much soy do Asians eat? A 1998 survey found that the average daily amount of soy protein consumed in Japan was about eight grams for men and seven for women -- less than two teaspoons.^[40] The famous Cornell China Study, conducted by Colin T. Campbell, found that legume consumption in China varied from 0 to 58 grams per day, with a mean of about twelve.^[41] Assuming that two-thirds of legume consumption is soy, then the maximum consumption is about 40 grams, or less than three tablespoons per day, with an average consumption of about nine grams, or less than two teaspoons. A survey conducted in the 1930s found that soy foods accounted for only 1.5 per cent of calories in the Chinese diet, compared with 65 per cent of calories from pork.^[42] (Asians traditionally cooked with lard,

not vegetable oil!)

Traditionally fermented soy products make a delicious, natural seasoning that may supply important nutritional factors in the Asian diet. But except in times of famine, Asians consume soy products only in small amounts, as condiments, and not as a replacement for animal foods -- with one exception. Celibate monks living in monasteries and leading a vegetarian lifestyle find soy foods quite helpful because they dampen libido.

It was a 1994 meta-analysis by Mark Messina, published in *Nutrition and Cancer*, that fuelled speculation on soy's anticarcinogenic properties.^[43] Messina noted that in 26 animal studies, 65 per cent reported protective effects from soy. He conveniently neglected to include at least one study in which soy feeding caused pancreatic cancer -- the 1985 study by Rackis.^[44] In the human studies he listed, the results were mixed. A few showed some protective effect, but most showed no correlation at all between soy consumption and cancer rates. He concluded that "the data in this review cannot be used as a basis for claiming that soy intake decreases cancer risk". Yet in his subsequent book, *The Simple Soybean and Your Health*, Messina makes just such a claim, recommending one cup or 230 grams of soy products per day in his "optimal" diet as a way to prevent cancer.

Thousands of women are now consuming soy in the belief that it protects them against breast cancer. Yet, in 1996, researchers found that women consuming soy protein isolate had an increased incidence of epithelial hyperplasia, a condition that presages malignancies.^[45] A year later, dietary genistein was found to stimulate breast cells to enter the cell cycle -- a discovery that led the study authors to conclude that women should not consume soy products to prevent breast cancer.^[46]

Phytoestrogens -- Panacea or Poison?

The male species of tropical birds carries the drab plumage of the female at birth and 'colours up' at maturity, somewhere between nine and 24 months.

In 1991, Richard and Valerie James, bird breeders in Whangerai, New Zealand, purchased a new kind of feed for their birds -- one based largely on soy protein.^[47] When soy-based feed was used, their birds 'coloured up' after just a few months. In fact, one bird-food manufacturer claimed that this early development was an advantage imparted by the feed. A 1992 ad for Roudybush feed formula showed a picture of the male crimson rosella, an Australian parrot that acquires beautiful red plumage at 18 to 24 months, already brightly coloured at 11 weeks old.

Unfortunately, in the ensuing years, there was decreased fertility in the birds, with precocious maturation, deformed, stunted and stillborn babies, and premature deaths, especially among females, with the result that the total population in the aviaries went into steady decline. The birds suffered beak and bone deformities, goitre, immune system disorders and pathological, aggressive behaviour. Autopsy revealed digestive organs in a state of disintegration. The list of problems corresponded with many of the problems the Jameses had encountered in their two children, who had been fed soy-based infant formula.

Startled, aghast, angry, the Jameses hired toxicologist Mike Fitzpatrick, PhD, to investigate further. Dr Fitzpatrick's literature review uncovered evidence that soy consumption has been linked to numerous disorders, including infertility, increased cancer and infantile leukaemia; and, in studies dating back to the 1950s,^[48] that genistein in soy causes endocrine disruption in animals. Dr Fitzpatrick also analysed the bird feed and found that it contained high levels of phytoestrogens, especially genistein. When the Jameses discontinued using soy-based feed, the flock gradually returned to normal breeding habits and behaviour.

The Jameses embarked on a private crusade to warn the public and government officials about toxins in soy foods, particularly the endocrine-disrupting isoflavones, genistein and diadzen. Protein Technology International received their material in 1994.

In 1991, Japanese researchers reported that consumption of as little as 30 grams or two tablespoons of soybeans per day for only one month resulted in a significant increase in thyroid-stimulating hormone.^[49] Diffuse goitre and hypothyroidism appeared in some of the subjects and many complained of constipation, fatigue and lethargy, even though their intake of iodine was adequate. In 1997, researchers from the FDA's National Center for Toxicological Research made the embarrassing discovery that the goitrogenic components of soy were the very same isoflavones.^[50]

Twenty-five grams of soy protein isolate, the minimum amount PTI claimed to have cholesterol-lowering effects, contains from 50 to 70 mg of isoflavones. It took only 45 mg of isoflavones in premenopausal women to exert significant biological effects, including a reduction in hormones needed for adequate thyroid function. These effects lingered for three months after soy consumption was discontinued.^[51]

One hundred grams of soy protein -- the maximum suggested cholesterol-lowering dose, and the amount recommended by Protein Technologies International -- can contain almost 600 mg of isoflavones,^[52] an amount that is undeniably toxic. In 1992, the Swiss health service estimated that 100 grams of soy protein provided the oestrogenic equivalent of the Pill.^[53]

In vitro studies suggest that isoflavones inhibit synthesis of oestradiol and other steroid hormones.^[54] Reproductive problems, infertility, thyroid disease and liver disease due to dietary intake of isoflavones have been observed for several species of animals including mice, cheetah, quail, pigs, rats, sturgeon and sheep.^[55]

It is the isoflavones in soy that are said to have a favourable effect on postmenopausal symptoms, including hot flushes, and protection from osteoporosis. Quantification of discomfort from hot flushes is extremely subjective, and most studies show that control subjects report reduction in discomfort in amounts equal to subjects given soy.^[56]

The claim that soy prevents osteoporosis is extraordinary, given that soy foods block calcium and cause vitamin D deficiencies. If Asians indeed have lower rates of osteoporosis than Westerners, it is because their diet provides plenty of vitamin D from shrimp, lard and seafood, and plenty of calcium from bone broths. The reason that Westerners have such high rates of osteoporosis is because they have substituted soy oil for butter, which is a traditional source of vitamin D and other fat-soluble activators needed for calcium absorption.

Birth Control Pills for Babies

But it was the isoflavones in infant formula that gave the Jameses the most cause for concern. In 1998, investigators reported that the daily exposure of infants to isoflavones in soy infant formula is 6 to 11 times higher on a body-weight basis than the dose that has hormonal effects in adults consuming soy foods. Circulating concentrations of isoflavones in infants fed soy-based formula were 13,000 to 22,000 times higher than plasma oestradiol concentrations in infants on cow's milk formula.^[57]

Approximately 25 per cent of bottle-fed children in the US receive soy-based formula -- a much higher percentage than in other parts of the Western world. Fitzpatrick estimated that an infant exclusively fed soy formula receives the oestrogenic equivalent (based on body weight) of at least five birth control pills per day.^[58] By contrast, almost no phytoestrogens have been detected in dairy-based infant formula or in human milk, even when the mother consumes soy products.

Scientists have known for years that soy-based formula can cause thyroid problems in babies. But what are the effects of soy products on the hormonal development of the infant, both male and female?

Male infants undergo a "testosterone surge" during the first few months of life, when testosterone levels may be as high as those of an adult male. During this period, the infant is programmed to express male characteristics after puberty, not only in the development of his sexual organs and other masculine physical traits, but also in setting patterns in the brain characteristic of male behaviour. In monkeys, deficiency of male hormones impairs the development of spatial perception (which, in humans, is normally more acute in men than in women), of learning ability and of visual discrimination tasks (such as would be required for reading).^[59] It goes without saying that future patterns of sexual orientation may also be influenced by the early hormonal environment. Male children exposed during gestation to diethylstilbestrol (DES), a synthetic oestrogen that has effects on animals similar to those of phytoestrogens from soy, had testes smaller than normal on maturation.^[60]

Learning disabilities, especially in male children, have reached epidemic proportions. Soy infant feeding -- which began in earnest in the early 1970s -- cannot be ignored as a probable cause for these tragic developments.

As for girls, an alarming number are entering puberty much earlier than normal, according to a recent study reported in the journal *Pediatrics*.^[61] Investigators found that one per cent of all girls now show signs of puberty, such as breast development or pubic hair, before the age of three; by age eight, 14.7 per cent of white girls and almost 50 per cent of African-American girls have one or both of these characteristics.

New data indicate that environmental oestrogens such as PCBs and DDE (a breakdown product of DDT) may cause early sexual development in girls.^[62] In the 1986 Puerto Rico Premature Thelarche study, the most significant dietary association with premature sexual development was not chicken -- as reported in the press -- but soy infant formula.^[63] The Woman, Infants and Children (WIC) program, which supplies free infant formula to welfare mothers, stresses soy formula for African Americans because they are supposedly allergic to

milk.

The consequences of this truncated childhood are tragic. Young girls with mature bodies must cope with feelings and urges that most children are not well-equipped to handle. And early maturation in girls is frequently a harbinger for problems with the reproductive system later in life, including failure to menstruate, infertility and breast cancer.

Parents who have contacted the Jameses recount other problems associated with children of both sexes who were fed soy-based formula, including extreme emotional behaviour, asthma, immune system problems, pituitary insufficiency, thyroid disorders and irritable bowel syndrome -- the same endocrine and digestive havoc that afflicted the Jameses' parrots.

Dissension in the Ranks

Organisers of the Third International Soy Symposium would be hard-pressed to call the conference an unqualified success. On the second day of the symposium, the London-based Food Commission and the Weston A. Price Foundation of Washington, DC, held a joint press conference, in the same hotel as the symposium, to present concerns about soy infant formula. Industry representatives sat stony-faced through the recitation of potential dangers and a plea from concerned scientists and parents to pull soy-based infant formula from the market. Under pressure from the Jameses, the New Zealand Government had issued a health warning about soy infant formula in 1998; it was time for the American government to do the same.

On the last day of the symposium, presentations on new findings related to toxicity sent a well-oxygenated chill through the giddy helium hype. Dr Lon White reported on a study of Japanese Americans living in Hawaii, that showed a significant statistical relationship between two or more servings of tofu a week and "accelerated brain aging".^[64] Those participants who consumed tofu in mid-life had lower cognitive function in late life and a greater incidence of Alzheimer's disease and dementia. "What's more," said Dr White, "those who ate a lot of tofu, by the time they were 75 or 80 looked five years older".^[65] White and his colleagues blamed the negative effects on isoflavones -- a finding that supports an earlier study in which postmenopausal women with higher levels of circulating oestrogen experienced greater cognitive decline.^[66]

Scientists Daniel Sheehan and Daniel Doerge, from the National Center for Toxicological Research, ruined PTI's day by presenting findings from rat feeding studies, indicating that genistein in soy foods causes irreversible damage to enzymes that synthesise thyroid hormones.^[67] "The association between soybean consumption and goiter in animals and humans has a long history," wrote Dr Doerge. "Current evidence for the beneficial effects of soy requires a full understanding of potential adverse effects as well."

Dr Claude Hughes reported that rats born to mothers that were fed genistein had decreased birth weights compared to controls, and onset of puberty occurred earlier in male offspring.^[68] His research suggested that the effects observed in rats ". . . will be at least somewhat predictive of what occurs in humans. There is no reason to assume that there will be gross malformations of fetuses but there may be subtle changes, such as neurobehavioral

attributes, immune function and sex hormone levels." The results, he said, "could be nothing or could be something of great concern . . . if mom is eating something that can act like sex hormones, it is logical to wonder if that could change the baby's development".^[69]

A study of babies born to vegetarian mothers, published in January 2000, indicated just what those changes in baby's development might be. Mothers who ate a vegetarian diet during pregnancy had a fivefold greater risk of delivering a boy with hypospadias, a birth defect of the penis.^[70] The authors of the study suggested that the cause was greater exposure to phytoestrogens in soy foods popular with vegetarians. Problems with female offspring of vegetarian mothers are more likely to show up later in life. While soy's oestrogenic effect is less than that of diethylstilbestrol (DES), the dose is likely to be higher because it's consumed as a food, not taken as a drug. Daughters of women who took DES during pregnancy suffered from infertility and cancer when they reached their twenties.

GRAS Status

Lurking in the background of industry hype for soy is the nagging question of whether it's even legal to add soy protein isolate to food. All food additives not in common use prior to 1958, including casein protein from milk, must have GRAS (Generally Recognized As Safe) status. In 1972, the Nixon administration directed a re-examination of substances believed to be GRAS, in the light of any scientific information then available. This re-examination included casein protein which became codified as GRAS in 1978. In 1974, the FDA obtained a literature review of soy protein because, as soy protein had not been used in food until 1959 and was not even in common use in the early 1970s, it was not eligible to have its GRAS status grandfathered under the provisions of the Food, Drug and Cosmetic Act.^[71]

The scientific literature up to 1974 recognised many antinutrients in factory-made soy protein, including trypsin inhibitors, phytic acid and genistein. But the FDA literature review dismissed discussion of adverse impacts, with the statement that it was important for "adequate processing" to remove them. Genistein could be removed with an alcohol wash, but it was an expensive procedure that processors avoided. Later studies determined that trypsin inhibitor content could be removed only with long periods of heat and pressure, but the FDA has imposed no requirements for manufacturers to do so.

The FDA was more concerned with toxins formed during processing, specifically nitrites and lysinoalanine.^[72] Even at low levels of consumption -- averaging one-third of a gram per day at the time -- the presence of these carcinogens was considered too great a threat to public health to allow GRAS status.

Soy protein did have approval for use as a binder in cardboard boxes, and this approval was allowed to continue, as researchers considered that migration of nitrites from the box into the food contents would be too small to constitute a cancer risk. FDA officials called for safety specifications and monitoring procedures before granting of GRAS status for food. These were never performed. To this day, use of soy protein is codified as GRAS only for this limited industrial use as a cardboard binder.

This means that soy protein must be subject to premarket approval procedures each time

manufacturers intend to use it as a food or add it to a food. Soy protein was introduced into infant formula in the early 1960s. It was a new product with no history of any use at all. As soy protein did not have GRAS status, premarket approval was required. This was not and still has not been granted. The key ingredient of soy infant formula is not recognised as safe.

The Next Asbestos?

"Against the backdrop of widespread praise . . . there is growing suspicion that soy -- despite its undisputed benefits -- may pose some health hazards," writes Marian Burros, a leading food writer for the *New York Times*. More than any other writer, Ms Burros's endorsement of a low-fat, largely vegetarian diet has herded Americans into supermarket aisles featuring soy foods. Yet her January 26, 2000 article, "Doubts Cloud Rosy News on Soy", contains the following alarming statement: "Not one of the 18 scientists interviewed for this column was willing to say that taking isoflavones was risk free." Ms Burros did not enumerate the risks, nor did she mention that the recommended 25 daily grams of soy protein contain enough isoflavones to cause problems in sensitive individuals, but it was evident that the industry had recognised the need to cover itself.

Because the industry is extremely exposed, contingency lawyers will soon discover that the number of potential plaintiffs can be counted in the millions and the pockets are very, very deep. Juries will hear something like the following: "The industry has known for years that soy contains many toxins. At first they told the public that the toxins were removed by processing. When it became apparent that processing could not get rid of them, they claimed that these substances were beneficial. Your government granted a health claim to a substance that is poisonous, and the industry lied to the public to sell more soy."

The "industry" includes merchants, manufacturers, scientists, publicists, bureaucrats, former bond financiers, food writers, vitamin companies and retail stores. Farmers will probably escape because they were duped like the rest of us. But they need to find something else to grow before the soy bubble bursts and the market collapses: grass-fed livestock, designer vegetables . . . or hemp to make paper for thousands and thousands of legal briefs.

References

1. Program for the Third International Symposium on the Role of Soy in Preventing and Treating Chronic Disease, Sunday, October 31, through Wednesday, November 3, 1999, Omni Shoreham Hotel, Washington, DC.
2. Houghton, Dean, "Healthful Harvest", *The Furrow*, January 2000, pp. 10-13.
3. Coleman, Richard J., "Vegetable Protein -- A Delayed Birth?" *Journal of the American Oil Chemists' Society* 52:238A, April 1975.
4. See www.unitedsoybean.org.
5. These are listed in www.soyonlineservice.co.nz.
6. *Wall Street Journal*, October 27, 1995.

7. Smith, James F., "Healthier tortillas could lead to healthier Mexico", *Denver Post*, August 22, 1999, p. 26A.
8. "Bakery says new loaf can help reduce hot flushes", *Reuters*, September 15, 1997.
9. "Beefing Up Burgers with Soy Products at School", *Nutrition Week*, Community Nutrition Institute, Washington, DC, June 5, 1998, p. 2.
10. Urquhart, John, "A Health Food Hits Big Time", *Wall Street Journal*, August 3, 1999, p. B1
11. "Soyabean Milk Plant in Kenya", *Africa News Service*, September 1998.
12. Simoons, Frederick J., *Food in China: A Cultural and Historical Inquiry*, CRC Press, Boca Raton, 1991, p. 64.
13. Katz, Solomon H., "Food and Biocultural Evolution: A Model for the Investigation of Modern Nutritional Problems", *Nutritional Anthropology*, Alan R. Liss Inc., 1987, p. 50.
14. Rackis, Joseph J. *et al.*, "The USDA trypsin inhibitor study. I. Background, objectives and procedural details", *Qualification of Plant Foods in Human Nutrition*, vol. 35, 1985.
15. Van Rensburg *et al.*, "Nutritional status of African populations predisposed to esophageal cancer", *Nutrition and Cancer*, vol. 4, 1983, pp. 206-216; Moser, P.B. *et al.*, "Copper, iron, zinc and selenium dietary intake and status of Nepalese lactating women and their breastfed infants", *American Journal of Clinical Nutrition* 47:729-734, April 1988; Harland, B.F. *et al.*, "Nutritional status and phytate: zinc and phytate X calcium: zinc dietary molar ratios of lacto-ovovegetarian Trappist monks: 10 years later", *Journal of the American Dietetic Association* 88:1562-1566, December 1988.
16. El Tiney, A.H., "Proximate Composition and Mineral and Phytate Contents of Legumes Grown in Sudan", *Journal of Food Composition and Analysis* (1989) 2:6778.
17. Ologhobo, A.D. *et al.*, "Distribution of phosphorus and phytate in some Nigerian varieties of legumes and some effects of processing", *Journal of Food Science* 49(1):199-201, January/February 1984.
18. Sandstrom, B. *et al.*, "Effect of protein level and protein source on zinc absorption in humans", *Journal of Nutrition* 119(1):48-53, January 1989; Tait, Susan *et al.*, "The availability of minerals in food, with particular reference to iron", *Journal of Research in Society and Health* 103(2):74-77, April 1983.
19. Phytate reduction of zinc absorption has been demonstrated in numerous studies. These results are summarised in Leviton, Richard, *Tofu, Tempeh, Miso and Other Soyfoods: The 'Food of the Future' -- How to Enjoy Its Spectacular Health Benefits*, Keats Publishing, Inc., New Canaan, CT, USA, 1982, p. 1415.
20. Mellanby, Edward, "Experimental rickets: The effect of cereals and their interaction with other factors of diet and environment in producing rickets", *Journal of the Medical Research Council* 93:265, March 1925; Wills, M.R. *et al.*, "Phytic Acid and Nutritional Rickets in Immigrants", *The Lancet*, April 8, 1972, pp. 771-773.
21. Rackis *et al.*, *ibid.*
22. Rackis *et al.*, *ibid.*, p. 232.
23. Wallace, G.M., "Studies on the Processing and Properties of Soymilk", *Journal of Science and Food Agriculture* 22:526-535, October 1971.
24. Rackis, *et al.*, *ibid.*, p. 22; "Evaluation of the Health Aspects of Soy Protein Isolates as Food Ingredients", prepared for FDA by Life Sciences Research Office, Federation of American Societies for Experimental Biology (9650 Rockville Pike, Bethesda, MD 20014), USA, Contract No. FDA

223-75-2004, 1979.

25. See www.truthinlabeling.org.
26. Rackis, Joseph, J., "Biological and Physiological Factors in Soybeans", *Journal of the American Oil Chemists' Society* 51:161A-170A, January 1974.
27. Rackis, Joseph J. *et al.*, "The USDA trypsin inhibitor study", *ibid.*
28. Torum, Benjamin, "Nutritional Quality of Soybean Protein Isolates: Studies in Children of Preschool Age", in *Soy Protein and Human Nutrition*, Harold L Wilcke *et al.* (eds), Academic Press, New York, 1979.
29. Zreik, Marwin, CCN, "The Great Soy Protein Awakening", *Total Health* 32(1), February 2000.
30. IEH Assessment on Phytoestrogens in the Human Diet, Final Report to the Ministry of Agriculture, Fisheries and Food, UK, November 1997, p. 11.
31. Food Labeling: Health Claims: Soy Protein and Coronary Heart Disease, Food and Drug Administration 21 CFR, Part 101 (Docket No. 98P-0683).
32. Sheegan, Daniel M. and Daniel R Doerge, Letter to Dockets Management Branch (HFA-305), February 18, 1999.
33. Anderson, James W. *et al.*, "Meta-analysis of the Effects of Soy Protein Intake on Serum Lipids", *New England Journal of Medicine* (1995) 333:(5):276-282.
34. Guy, Camille, "Doctors warned against magic, quackery", *New Zealand Herald*, September 9, 1995, section 8, p. 5.
35. Sander, Kate and Hilary Wilson, "FDA approves new health claim for soy, but little fallout expected for dairy", *Cheese Market News*, October 22, 1999, p. 24.
36. Enig, Mary G. and Sally Fallon, "The Oiling of America", NEXUS Magazine, December 1998-January 1999 and February-March 1999; also available at www.westonaprice.org/facts_about_fats/oiling.html.
37. *Natural Medicine News* (L & H Vitamins, 32-33 47th Avenue, Long Island City, NY 11101), USA, January/February 2000, p. 8.
38. Harras, Angela (ed.), *Cancer Rates and Risks*, National Institutes of Health, National Cancer Institute, 1996, 4th edition.
39. Searle, Charles E. (ed.), *Chemical Carcinogens*, ACS Monograph 173, American Chemical Society, Washington, DC, 1976.
40. Nagata, C. *et al.*, *Journal of Nutrition* (1998) 128:209-213.
41. Campbell, Colin T. *et al.*, *The Cornell Project in China*.
42. Chang, K.C. (ed.), *Food in Chinese Culture: Anthropological and Historical Perspectives*, New Haven, 1977.
43. Messina, Mark J. *et al.*, "Soy Intake and Cancer Risk: A Review of the In Vitro and In Vivo Data", *Nutrition and Cancer* (1994) 21(2):113-131.
44. Rackis *et al.*, "The USDA trypsin inhibitor study", *ibid.*
45. Petrakis, N.L. *et al.*, "Stimulatory influence of soy protein isolate on breast secretion in pre- and

post-menopausal women", *Cancer Epid. Bio. Prev.* (1996) 5:785-794.

46. Dees, C. *et al.*, "Dietary estrogens stimulate human breast cells to enter the cell cycle", *Environmental Health Perspectives* (1997) 105(Suppl. 3):633-636.
47. Woodhams, D.J., "Phytoestrogens and parrots: The anatomy of an investigation", *Proceedings of the Nutrition Society of New Zealand* (1995) 20:22-30.
48. Matrone, G. *et al.*, "Effect of Genistin on Growth and Development of the Male Mouse", *Journal of Nutrition* (1956) 235-240.
49. Ishizuki, Y. *et al.*, "The effects on the thyroid gland of soybeans administered experimentally in healthy subjects", *Nippon Naibunpi Gakkai Zasshi* (1991) 767:622-629.
50. Divi, R.L. *et al.*, "Anti-thyroid isoflavones from the soybean", *Biochemical Pharmacology* (1997) 54:1087-1096.
51. Cassidy, A. *et al.*, "Biological Effects of a Diet of Soy Protein Rich in Isoflavones on the Menstrual Cycle of Premenopausal Women", *American Journal of Clinical Nutrition* (1994) 60:333-340.
52. Murphy, P.A., "Phytoestrogen Content of Processed Soybean Foods", *Food Technology*, January 1982, pp. 60-64.
53. Bulletin de L'Office Fédéral de la Santé Publique, no. 28, July 20, 1992.
54. Keung, W.M., "Dietary oestrogenic isoflavones are potent inhibitors of B-hydroxysteroid dehydrogenase of *P. testosteronei*", *Biochemical and Biophysical Research Committee* (1995) 215:1137-1144; Makela, S.I. *et al.*, "Estrogen-specific 12 B-hydroxysteroid oxidoreductase type 1 (E.C. 1.1.1.62) as a possible target for the action of phytoestrogens", *PSEBM* (1995) 208:51-59.
55. Setchell, K.D.R. *et al.*, "Dietary oestrogens -- a probable cause of infertility and liver disease in captive cheetahs", *Gastroenterology* (1987) 93:225-233; Leopald, A.S., "Phytoestrogens: Adverse effects on reproduction in California Quail," *Science* (1976) 191:98-100; Drane, H.M. *et al.*, "Oestrogenic activity of soya-bean products", *Food, Cosmetics and Technology* (1980) 18:425-427; Kimura, S. *et al.*, "Development of malignant goiter by defatted soybean with iodine-free diet in rats", *Gann.* (1976) 67:763-765; Pelissero, C. *et al.*, "Oestrogenic effect of dietary soybean meal on vitellogenesis in cultured Siberian Sturgeon *Acipenser baeri*", *Gen. Comp. End.* (1991) 83:447-457; Braden *et al.*, "The oestrogenic activity and metabolism of certain isoflavones in sheep", *Australian J. Agricultural Research* (1967) 18:335-348.
56. Ginsburg, Jean and Giordana M. Prelevic, "Is there a proven place for phytoestrogens in the menopause?", *Climacteric* (1999) 2:75-78.
57. Setchell, K.D. *et al.*, "Isoflavone content of infant formulas and the metabolic fate of these early phytoestrogens in early life", *American Journal of Clinical Nutrition*, December 1998 Supplement, 1453S-1461S.
58. Irvine, C. *et al.*, "The Potential Adverse Effects of Soybean Phytoestrogens in Infant Feeding", *New Zealand Medical Journal* May 24, 1995, p. 318.
59. Hagger, C. and J. Bachevalier, "Visual habit formation in 3-month-old monkeys (*Macaca mulatta*): reversal of sex difference following neonatal manipulations of androgen", *Behavior and Brain Research* (1991) 45:57-63.
60. Ross, R.K. *et al.*, "Effect of in-utero exposure to diethylstilbestrol on age at onset of puberty and on post-pubertal hormone levels in boys", *Canadian Medical Association Journal* 128(10):1197-8, May 15, 1983.

61. Herman-Giddens, Marcia E. *et al.*, "Secondary Sexual Characteristics and Menses in Young Girls Seen in Office Practice: A Study from the Pediatric Research in Office Settings Network", *Pediatrics* 99(4):505-512, April 1997.
62. *Rachel's Environment & Health Weekly* 263, "The Wingspread Statement", Part 1, December 11, 1991; Colborn, Theo, Dianne Dumanoski and John Peterson Myers, *Our Stolen Future*, Little, Brown & Company, London, 1996.
63. Freni-Titulaer, L.W., "Premature Thelarch in Puerto Rico: A search for environmental factors", *American Journal of Diseases of Children* 140(12):1263-1267, December 1986.
64. White, Lon, "Association of High Midlife Tofu Consumption with Accelerated Brain Aging", Plenary Session #8: Cognitive Function, The Third International Soy Symposium, November 1999, Program, p. 26.
65. Altonn, Helen, "Too much tofu induces 'brain aging', study shows", *Honolulu Star-Bulletin*, November 19, 1999.
66. *Journal of the American Geriatric Society* (1998) 46:816-21.
67. Doerge, Daniel R., "Inactivation of Thyroid Peroxidase by Genistein and Daidzein in Vitro and in Vivo; Mechanism for Anti-Thyroid Activity of Soy", presented at the November 1999 Soy Symposium in Washington, DC, National Center for Toxicological Research, Jefferson, AR 72029, USA.
68. Hughes, Claude, Center for Women's Health and Department of Obstetrics & Gynecology, Cedars-Sinai Medical Center, Los Angeles, CA.
69. Soy Intake May Affect Fetus", *Reuters News Service*, November 5, 1999.
70. "Vegetarian diet in pregnancy linked to birth defect", *BJU International* 85:107-113, January 2000.
71. FDA ref 72/104, Report FDABF GRAS -- 258.
72. "Evaluation of the Health Aspects of Soy Protein Isolates as Food Ingredients", prepared for FDA by Life Sciences Research Office, Federation of American Societies for Experimental Biology (FASEB) (9650 Rockville Pike, Bethesda, MD 20014, USA), Contract No, FDA 223-75-2004, 1979.

About the Authors:

Sally Fallon is the author of *Nourishing Traditions: The Cookbook that Challenges Politically Correct Nutrition and the Diet Dictocrats* (1999, 2nd edition, New Trends Publishing, tel +1 877 707 1776 or +1 219 268 2601) and President of the Weston A. Price Foundation, Washington, DC (www.WestonAPrice.org).

Mary G. Enig, PhD, is the author of *Know Your Fats: The Complete Primer for Understanding the Nutrition of Fats, Oils and Cholesterol* (2000, Bethesda Press, www.BethesdaPress.com), is President of the Maryland Nutritionists Association and Vice President of the Weston A. Price Foundation, Washington, DC.

The authors wish to thank Mike Fitzpatrick, PhD, and Valerie and Richard James for their help in preparing this article.

Ironically, the soy dietâ€”including â€œmeatâ€ patties containing 60-70 percent soy protein, fake soy cheese, soy sausage and soy flour or soy protein added to most baked goodsâ€”replaced one that was quite healthy for the prisoners. Well into the late 1990s, the prisoners operated prison farms and abattoirs, producing most of their own food. Studies concerning the impacts of soy on human health and the effects of its cultivation on the social-environmental domain were reviewed to analyze the political context of the discussion surrounding soy and the socio-environmental repercussions of its cultivation. Based on the sociology of scientific knowledge and the environmental sociology, we identified a thin line between healthy and risky food, which is vulnerable to different reflexively constructed influences.Â Soy Alert -- Tragedy and Hype The Third International Soy Symposium. Article. Sally Fallon. Fallon S, Enig M. Soy Alert â€“ Tragedy and Hype. Nexus Magazine. 2000 Apr-May;7(3). 5. Solomons NW, Torun B. Infantile malnutrition in the tropics.Â **Tragedy and Hype: The Third International Soy Symposium â€“ Part II. Sally Fallon and Mary G. Enig, PhD Life Extension Magazine July 2010. Is Soy Safe?.