

Invertebrates of temporary waters in gnammas on granite outcrops in Western Australia

I A E Bayly

Department of Biological Sciences,
Monash University, Clayton VIC 3168

Abstract

Thirty-six flooded gnammas (rock pools), distributed between 17 different granite outcrops, were each sampled once only during the winter of 1990. The water of most pools was acidic (pH < 7.0) and of low conductivity ($K_{25} < 200 \mu\text{S cm}^{-1}$). A total of 88 invertebrate taxa were found with *Boeckella opaqua* (Copepoda; 19 occurrences), *Cyprretta baylyi* (Ostracoda; 17) and *Neothrix armata* (Cladocera; 14) most common. A high proportion of the Crustacea consisted of species endemic to Western Australia. Six new species of *Cypricercus* (Ostracoda) were discovered. The mean number of taxa was 8.2 per pool. There was a highly significant positive correlation between species richness and the logarithm of both the pool volume and pool area. The nature of the fauna is reviewed within the framework of the type of adaptations employed by the animals for tolerating or avoiding the dry phase.

Introduction

The word "gnamma" is of Aboriginal origin and refers to a depression that has been weathered out of the surface of an outcrop of bare rock; some overseas workers (e.g. Smith 1941) have referred to these depressions as "weather pits". Gnammas, or weather pits, are commonly found on granite outcrops and especially on the top of domed inselbergs. Despite the widespread occurrence of gnammas in Australia, the aquatic biota of rain-filled gnammas has received remarkably little attention. This neglect might be partly explicable on the basis that, because gnammas are decidedly at the small end of the size spectrum of lentic waters (pool-pond-lake), they have been regarded as being of little interest or consequence to aquatic ecologists. In fact, small size confers on gnammas a number of methodological advantages, such as ease of sampling and experimental manipulation, as subjects for an ecologist. Sheldon (1984), for example, draws attention to the good potential of rock pools for comparative and experimental studies of the aerial colonization dynamics of adult aquatic insects. This claim for gnammas, along perhaps with phytotelmata (pockets of water held in plants such as the Albany Pitcher Plant; Bayly 1984), as being quintessential aquatic microcosms, is appealing. It may be claimed with some confidence, therefore, that the potential of studies of gnammas for illustrating ecological principles is just as great, if not greater, than that of large lakes.

Among the earliest written observations on the gnammas of Western Australian granite outcrops are those of the explorer Carnegie (1898). Interesting comments on the form, distribution and likely origin of granitic gnammas in this State were also recorded by the geologist Jutson (1934). Taxonomic work on invertebrates collected from Western Australian

gnammas has been undertaken by Wolf (1911), Fairbridge (1945), Petkovski (1973), Wallwork (1981), De Deckker (1981), Frey (1991, 1998), Lansbury (1995), Smirnov & Bayly (1995) and Benzie & Bayly (1996). Studies of the ecology of these gnammas, or of specific animals living in them, have been made by Jones (1971, 1974) and Bayly (1982).

Thus far only a limited attempt has been made to deal with the fauna of these peculiar and distinctive aquatic habitats in a comprehensive manner. The aim of this paper is to provide a fairly complete account of the invertebrate inhabitants of gnammas located in the more southern regions of Western Australia. This is a necessary basis for the future realisation of the above-mentioned potential of gnammas for facilitating ecological studies. Ecological aspects of the present study include the documentation of some salient chemical features of gnamma waters and the examination of species richness in relation to pool size.

This account is based on collections made from 36 gnammas, distributed between 17 different granite outcrops, during the winter of 1990. Thirteen of the 17 outcrops were in the form of domical inselbergs while the remainder were small flat surfaces. With one exception, the gnammas were shallow, flat-bottomed pan-gnammas *sensu* Twidale & Corbin (1963). The exception was a deep pit-gnamma located on War Rock.

Materials and Methods

The maximum depth, length and width (the greatest width at right angles to the line of maximum length) were measured at each pool. Hydrogen ion concentration and conductivity of water samples collected in a polyethylene bottle were determined with a Metrohm E588 pH-meter and Radiometer CDM2e conductivity meter respectively. Invertebrates were collected with a rectangular-framed net conforming with that described by Bayly (1982), except that the mesh

aperture was 150 μm , and preserved in either 10% formalin or 90% ethanol.

The product of the maximum length and maximum width of a pool was used to approximate (consistently overestimate) the true areas of the pools. Likewise, the product of the maximum length, maximum width and maximum depth was used to approximate (consistently overestimate) the true volumes of the pools. The approximated areas and volumes of the pools were log-transformed for correlation calculations.

Results

Data on the location and physical dimensions of the 36 pools as well as their water chemistry are presented in Table 1. With one exception, pH was within the range 4.6–7.9, but for 31 of the pools it was less than 7.0. With three exceptions, the conductivity (K_{25}) was less than 1000 $\mu\text{S cm}^{-1}$, and for the 33 pools whose conductivity was less than this value the mean was 147 $\mu\text{S cm}^{-1}$.

The results of taxonomic studies are summarized in Table 2. A total of 88 taxa were recorded and the mean number was 8.2 per gnamma. Branchiopod crustaceans were found in 18 pools, with *Limnadia* the most frequently occurring taxon. Cladocera occurred in 26 pools, with *Neothrix armata* the commonest species. Copepoda were found in 23 pools, with *Boeckella opaqua* occurring most frequently. Ostracods occurred in all but one gnamma and were the only animals collected from four gnammas. The commonest ostracod was *Cyprretta baylyi*. At least six new species of *Cypricercus* were recorded. Insects were found in 19 pools, with *Anisops thienemanni* the commonest species. Pools 16 and 21 yielded a disproportionately high number of insect species. Oribatid mites were restricted to two small coastal pools.

There is a highly significant, positive correlation between species richness and the logarithm of approximated pool volume (Fig 1; Pearson $r = 0.595$, $n = 36$, $P < 0.001$). If one outlier (two taxa from pool 19 with log volume 4.78) is removed from the data set, then the

Table 1
Physico-chemical features of Western Australian granite rock-pools sampled June-August 1990.

Locality number and name	Location (lat S, long E)	Sampling date	Maximum depth (cm)	Max. length x max. width (m^2)	pH	Conductivity (K_{25} ; $\mu\text{S cm}^{-1}$)
1. Coragina Rock (a)	32° 55', 123° 30'	24.vi.1990	25	16 x 7	8.7	91
2. Coragina Rock (b)	" "	"	6	3 x 2	6.4	59
3. Mt Madden (a)	33° 14', 119° 50'	26.vi.1990	30	30 x 10	6.3	74
4. Mt Madden (b)	" "	"	10	12 x 2.5	6.0	96
5. Mt Madden (c)	" "	"	15	5 x 2.5	6.1	38
6. Mt Madden (d)	" "	"	20	7 x 5	6.0	61
7. Cable Beach (a)	35° 07', 117° 54'	27.vi.1990	4	1 x 0.3	7.1	1480
8. Cable Beach (b)	" "	"	2	1 x 0.3	7.0	1490
9. Frenchman Bay Rd (a)	35° 06', 117° 57'	27.vi.1990	2	3 x 2	7.3	380
10. Frenchman Bay Rd (b)	" "	"	5	0.8 x 0.2	6.4	457
11. Muirillup Rock (a)	34° 39', 116° 15'	2.vii.1990	12	3 x 1.5	4.6	93
12. Muirillup Rock (b)	34° 39', 116° 15'	"	6	2 x 1.5	5.5	101
13. Cape Leeuwin (a)	34° 22', 115° 08'	6.vii.1990	2	2.5 x 1.2	6.7	450
14. Cape Leeuwin (b)	" "	"	4	5 x 1.5	6.5	8040
15. Bilya Rock	29° 00', 115° 51'	12.vii.1990	11	3 x 2.5	6.1	90
16. War Rock (a)	29° 05', 116° 00'	13.vii.1990	22	25 x 9	5.2	55
17. War Rock (b)	" "	"	>100	6 x 2	7.9	798
18. War Rock (c)	" "	"	12	4 x 2	6.1	38
19. Bunjil Rock (a)	29° 39', 116° 21'	13.vii.1990	100	10 x 6	6.8	60
20. Bunjil Rock (b)	" "	"	9	8 x 5	6.1	34
21. Petrudor Rocks (a)	30° 25', 116° 58'	14.viii.1990	80	11 x 11	6.5	116
22. Petrudor Rocks (b)	" "	"	6	3 x 1.5	6.0	186
23. Petrudor Rocks (c)	" "	"	19	6 x 3	6.2	190
24. Elachbutting (a)	30° 36', 118° 37'	15.viii.1990	10	6 x 2	5.1	164
25. Elachbutting (b)	" "	"	19	6 x 5	5.7	162
26. Sanford Rock (a)	31° 14', 118° 46'	15.viii.1990	26	8 x 3	6.3	102
27. Sanford Rock (b)	" "	"	12	2 x 2	6.7	88
28. Jilbadgie Rocks (a)	31° 29', 119° 13'	16.viii.1990	18	6 x 4	6.2	34
29. Jilbadgie Rocks (b)	" "	"	20	13 x 6	6.3	42
30. Mt Hampton (a)	31° 45', 119° 04'	16.viii.1990	33	10 x 9	6.3	144
31. Mt Hampton (b)	" "	"	13	4 x 3	6.4	64
32. Yorkrakine Rock (a)	31° 26', 117° 31'	26.viii.1990	9	7 x 5	6.4	113
33. Yorkrakine Rock (b)	" "	"	12	4 x 3	6.2	146
34. King Rocks (a)	32° 19', 119° 09'	28.viii.1990	20	14 x 10	6.6	105
35. King Rocks (b)	" "	"	30	16 x 8	6.4	91
36. Wave Rock	32° 27', 118° 54'	28.viii.1990	12	9 x 3	6.3	113

Table 2

List of taxa and their occurrences.

Taxa	Pool (see Table 1)	Total number of occurrences	Taxa	Pool (see Table 1)	Total number of occurrences
CRUSTACEA: ANOSTRACA					
<i>Branchinella longirostris</i> Wolf	5,24,25,26,27,31	6	<i>Cypricercus</i> sp or spp	6,20,22,26,33,34	6
<i>B. sp</i>	34	1	<i>C. n sp 1</i>	2,3	2
CRUSTACEA: CONCHOSTRACA					
<i>Cyzicus</i> sp	4,5,6,26	4	<i>C. n sp 2</i>	3,4,5,8,30	5
<i>Limnadia</i> sp or spp	15,22,25,27,28,29,30, 31,33,36	10	<i>C. n sp 3</i>	3,34	2
<i>Lynceus</i> sp	17,23	2	<i>C. n sp 4</i>	6,9,13,14,15,16	6
CRUSTACEA: CLADOCERA					
Chydoridae					
<i>Alona cf. setuloides</i>	32,33	2	<i>C. n sp 5</i>	30	1
<i>A. n sp</i>	33	1	<i>C. n sp 6</i>	36	1
<i>Alonella - excisa</i> group	11	1	<i>Heterocypris incongruens</i> (Rahmdor)	2	1
<i>Biapertura - macropa</i> group			<i>Ilydromus amplicolis</i> De Deckker	2,10,18,20,23,29,31,32,33	9
sp1	15	1	<i>I. candonites</i> De Deckker	9,12,16,18	4
sp2	11,35	2	<i>I. sp</i>	4,15,21,24,27,36	6
sp3	5,12,24,25,26,29,30,34,36	9	<i>Limnocythere mowbrayensis</i> Chapman	2,3,7,34,35,36	6
<i>B. rigidicaudis</i> Smirnov	6,27,28	3	<i>L. sp</i>	8	1
<i>B. n sp</i>	9,11	2	<i>L. n sp?</i>	13	1
<i>Ephemeroporus-barroisi</i> group			<i>L. n sp</i>	32	1
sp1	6,34,35	4	<i>Sarscypridopsis aculaeata</i> (Costa)	17,21,23	3
sp2	11	1	ACARI: ORIBATEI		
<i>Monospilus diporus</i> Smirnov & Timms	4,23,26	3	<i>Scapheremaeus</i> sp	78	2
<i>Planicirculus alticarinatus</i> Frey	26,31	2	<i>Trimalaconothrus</i> sp	78	2
<i>Plurispina chauliodus</i> Frey	12	1	INSECTA: COLEOPTERA		
<i>P. multituberculata</i> Frey	2,6,25,33	4	Dytiscidae		
<i>Rak stagnensis</i> Frey	1,4,5,27,29,34	6	<i>Allodessus bistrigatus</i> (Clark)	16,21,25	3
<i>Leberis aenigmatica</i> Smirnov	27,29,31,34	4	<i>Eretes australis</i> (Erichson)	21	1
Other families					
<i>Daphnia jollyi</i> Petkovski	23,26,29,31	4	<i>Lanceotes lanceolatus</i> (Clark)	6	1
<i>Neothrix armata</i> Gurney	1,3,6,12,15,18,21, 23,24,26,30,34,35,36	14	<i>Megaporus howitti</i> (Clark)	3,16	2
<i>Macrothrix breviseta</i> Smirnov	30,35	2	<i>Necterosoma darwini</i> (Babington)	11	1
<i>M. hardingi</i> Petkovski	24,27	2	<i>Paroster niger</i> Watts	78	2
<i>M. indistincta</i> Smirnov	1,25	2	<i>Sternopriscus multimaculatus</i> (Clark)	16,25	2
<i>M. longiseta</i> Smirnov	2	1	Hydrophilidae		
<i>Ceriodaphnia</i> sp	1,26,27,30,34,35,36	7	<i>Berosus approximatus</i> Fairmaire	16	1
<i>Moina</i> sp	1,2,15,18	4	<i>Limnoxenus zelandicus</i> (Broun)	6	1
CRUSTACEA: COPEPODA					
Calanoida					
<i>Boeckella opaqua</i> Fairbridge	1,2,3,4,5,6,24,25,26,27,28, 29,30,31,32,33,34,35,36	19	INSECTA: DIPTERA		
<i>B. triarticulata</i> (Thomson)	17,19,21,23	4	Ceratopogonidae		
<i>Calamoecia ampulla</i> (Searle)	21	1	<i>Dasyhelea</i> sp	2,22,25	3
Cyclopoida					
<i>Mesocyclops australiensis</i> (Sars)	21	1	Chironomidae		
<i>Metacyclops</i> sp	3,34	2	<i>Ablabesmyia</i> sp	28	1
<i>Microcyclops varicans</i> (Sars)	1,21	2	<i>Allostrissocladus</i> sp	28,30	2
Harpacticoida					
	11	1	<i>Chironomus tepperi</i> Skuse	2,23	2
CRUSTACEA: OSTRACODA					
<i>Alboa n sp</i>	16	1	<i>Dicrotendipes</i> sp	11	1
<i>Bennelongia barangaroo</i> De Deckker	1,5,23,25,29	5	<i>Paraborniola</i> sp	16	1
<i>B. sp</i>	3,21	2	<i>?Paratendipes</i> sp	33	1
<i>Candonocypris novaezelandiae</i> (Baird)	16	1	Orthoclaadiinae		
<i>Cyprretta baylyi</i> McKenzie	3,4,6,11,12,16,18,20,22,23, 25,26,27,28,32,33,36	17		25,28,30	3
<i>C. sp</i>	2,10,15,22,25,28,31, 32,33,34,35	11	INSECTA: HEMIPTERA		
			Corixidae		
			<i>Agraptocorixa parvipunctata</i> (Hale)	3,6,16,21,23	5
			<i>Diaprepocoris personata</i> Hale	11	1
			<i>Micronecta annae</i> Kirkaldy	16,21	2
			<i>M. gracilis</i> Hale	21	1
			<i>M. robusta</i> Hale	4,6,25,26,36	5
			<i>Sigara mullaka</i> Lansbury	26	1
			Notonectidae		
			<i>Anisops baylyi</i> Lansbury	5,6,16	3
			<i>A. gratus</i> Hale	21	1
			<i>A. hyperion</i> Kirkaldy	3,6,17,21	4
			<i>A. stali</i> Kirkaldy	17,21	2
			<i>A. thienemanni</i> Lundblad	3,6,16,17,21,23,25,30	8




Figure 1. Plot of species richness against the logarithm (base 10) of the estimated volume (in litres) for all 36 gnammas. There is a highly significant positive correlation between these two variables (Pearson $r = 0.595$, $n = 36$, $P < 0.001$). The outlier at two species and log volume 4.78 is pool number 19 in Table 1.

correlation is considerably improved ($r = 0.696$, $n = 35$, $P < 0.001$). There is also a highly significant positive correlation between species richness (Pearson $r = 0.577$, $n = 36$, $P < 0.001$) and the logarithm of approximated pool area (Pearson $r = 0.641$, $n = 35$, $P < 0.001$).

Discussion

With seven exceptions, the conductivity (K_{25}) of the pools (Table 1) was less than $200 \mu\text{S cm}^{-1}$ - a finding broadly consistent with that of Bayly (1982) for a series of 19 granite rock pools in Western Australia. Six of the seven exceptions were pools with a maritime location (Cable Beach, Frenchman Bay and Cape Leeuwin) and whose high conductivities were clearly caused by the accession of ions from marine salt spray. The remaining exception was a deep pit-gnamma located on War Rock; this gnamma probably rarely loses accumulated ions by means of overflow.

The pH of almost all of the inland pools is acidic, as expected for a granite substratum. The exceptionally high pH recorded for Coragina Rock (a; Table 1) was discussed by Bayly (1992) who attributed it to the use of cement mortar between the exfoliated slabs of granite used to raise the level of the pool.

Although no quantitative data are available, it is noteworthy that granite rock-pools lack the high turbidity that is so characteristic of pools lying on the regolith, especially in arid regions.

The total number of taxa listed in Table 2 is 88. This, however, is an underestimate of the total number of metazoan invertebrate taxa because black planarians (*Bothrosostoma?*) and nematodes were present in several collections but were not identified and are omitted from Table 2. The total of 88 taxa greatly exceeds the 18 invertebrate taxa listed by Bishop (1974) as occurring in a series of shallow pools located on sandstone at Kanangra Walls in the Blue Mountains, New South Wales. The 88 taxa from all 36 pools and the 17 taxa from the richest pool (number 21) may also be compared with the 121 invertebrate taxa reported by Lake *et al.* (1989) from a single temporary pond (not located on a granitic substratum) in western Victoria that was repeatedly sampled over 7 months.

The animals listed in Table 2 may be classified into four groups according to their adaptations for tolerating or avoiding the dry phase following the system of Wiggins *et al.* (1980) as modified by Williams (1985).

- **Group A** (Group I of Wiggins *et al.* 1980) consists of permanent residents of the pools that are capable of only passive dispersal. These organisms are dormant during the dry season and typically avoid desiccation as a resistant stage (shelled egg or embryo). All of the Crustacea listed in Table 2 should be assigned to this group, a salient feature of which is the high proportion of species endemic to Western Australia. Such endemics include *Branchinella longirostris*, *Daphnia jollyi*, *Macrothrix hardingi*, *Boeckella opaqua*, *Ilyodromus amplicolis*, *I. candonites*, *Leberis aenigmatosa*, *Plurispina*

chauliodus and *P. multituberculata*. This list does not include the new species of *Cypricercus* some of which are likely to prove endemic. A distinctive negative feature of this group is the absence of notostracans. Despite their possession of a suite of adaptations (including resistant eggs) for occupancy of small temporary waters, an unknown factor excluded notostracans from granite rock pools studied here. A possible reason is that the concentration of calcium in waters on granite is so low that it is physiologically impossible for *Lepidurus* or *Triops* to fabricate their relatively large carapace. The carapaces of *Ilyodromus* and other ostracod taxa living in granite rock-pools contain practically no calcium; instead their carapaces are strengthened by non-calcareous ridges to compensate for this deficiency P De Deckker (*pers comm*).

- **Group B** (Group II of Wiggins *et al.* 1980) comprises animals (typically insects) that are capable of active dispersal but which are typically still present during the dry season in a dormant form (quiescent larvae). *Allotrissocladus*, *Paraborniola* and *Dasyhelea* are assigned to this group. The ability of the larvae of these taxa to resist desiccation in gnammas was investigated by Jones (1971, 1975).

- **Group C** (Group IV of Wiggins *et al.* 1980) consists of animals (typically insects) capable of active dispersal and which have a discontinuous presence in the pools; they avoid the dry period altogether by emigrating to permanent waters before the pools dry up. To this group are assigned all the Dystiscidae, Hydrophilidae, Corixidae and Notonectidae listed in Table 2, some 20 species in all.

- **Group D** [Williams' (1985) replacement for Group III of Wiggins *et al.*] comprises animals which lack a resistant stage in their life cycle but possess exceptionally good dispersal ability thus allowing a discontinuous presence in the confines of a temporary water basin. A single species, *Chironomus tepperi*, may be assigned to this group. The ability of this species to rapidly colonize temporary waters has been demonstrated by Maher & Carpenter (1984).

The biology of the oribatid mites, *Chudalupia meridionalis* (see Bayly 1992), *Scapheremaeus* sp and *Trimalaconothrus* sp is insufficiently known to allow their assignment within the Wiggins-Williams system.

Wiggins *et al.* (1980) mentioned the habit of burrowing into bottom sediments as a behavioural adaptation for avoiding desiccation in temporary pools. While this pattern of behaviour may provide an escape for some animals in most pools, there is little scope for it in most pan-gnammas because of minimal amounts of sediment and the impossibility of burrowing into granite. Deep pit-gnammas left undisturbed for long periods may accumulate a significant amount of sediment and provide an exception.

Fryer (1985), on the basis of studies in England, concluded that chydorid cladocerans "show an unambiguous preference for large water bodies". However, this conclusion is of doubtful validity in the Australian context (Frey 1998). Although sampled once only, some gnammas contained four species of Chydoridae and there is presently no evidence that this number is significantly exceeded in large Australian

lakes. Timms (1981), for example, sampled the littoral region of Lake Purrumbete, a large (552 ha) freshwater lake in Western Victoria, monthly for a year, but recorded only six chydorid species.

The finding of a significant positive correlation between species richness and the volume of the pools is consistent with earlier investigations. Ranta (1982) reported a significant correlation between the number of water beetle species and the logarithm of pool volume in a series of 20 coastal (marine supralittoral) rock pools located in the Baltic region. A similar finding for a greater diversity of taxa was made by March & Bass (1995) for a series of six temporary pools located in Oklahoma.

Acknowledgements: I am very grateful to the following for providing taxonomic identifications; the late Professor D G Frey (Chydoridae), P De Deckker (Ostracoda), I Lansbury (Corixidae and Notonectidae), M C Geddes (Anostraca and Conchostraca), D W Morton (Cyclopoida), M Colloff (Oribatei), D H D Edward (Chironomidae and Ceratopogonidae), J F Lawrence (Hydrophilidae) and C H S Watts (Dytiscidae). Because of the importance of ostracods in gnammas, P De Deckker's task was a particularly onerous one. I thank the Royal Society of Western Australia for inviting me to contribute to the Symposium and for paying my travelling expenses. I am indebted to L Dembinski for the word processing.

References

- Bayly I A E 1982 Invertebrate fauna and ecology of temporary pools on granite outcrops in southern Western Australia. *Australian Journal of Marine and Freshwater Research* 33:599-606.
- Bayly I A E 1984 Phytotelmata: terrestrial plants as hosts for aquatic insect communities [Book review] *Australian Journal of Ecology* 9:164-165
- Bayly I A E 1992 Freshwater havens. *Landscape* 7(4):49-53.
- Benzie J A H & Bayly I A E 1996 Male and epihippial female *Daphnia jollyi* Petkovski, 1973 discovered in Western Australia and the parthenogenetic female redescribed. *Hydrobiologia* 331:171-181.
- Bishop J A 1974 The fauna of temporary rain pools in eastern New South Wales. *Hydrobiologia* 44:319-323.
- Carnegie D W 1898 *Spinifex* and *Sand*. Arthur Pearson, London.
- De Deckker P 1981 Ostracoda from Australian inland waters - notes on taxonomy and ecology. *Proceedings of the Royal Society of Victoria* 93:43-85.
- Fairbridge W S 1945 West Australian freshwater calanoids (Copepoda). I Three new species of *Boeckella*, with a description of the developmental stages of *B. opaquana* n sp and a key to the genus. *Journal of the Royal Society of Western Australia* 29:25-65.
- Frey D G 1991 The species of *Pleuroxus* and of three related genera (Anomopoda, Chydoridae) in southern Australia and New Zealand. *Records of the Australian Museum* 43:291-372.
- Frey D G 1998 Expanded description of *Leberis aenigmatica* Smirnov (Anomopoda: Chydoridae): further indication of the biological isolation between western and eastern Australia. *Hydrobiologia* (in press).
- Fryer G 1985 Crustacean diversity in relation to the size of water bodies: some facts and problems. *Freshwater Biology* 15:347-361.
- Jones R E 1971 The ecology of some species of Diptera on granite outcrops. PhD Thesis. University of Western Australia, Perth.
- Jones R E 1974 The effects of size-selective predation and environmental variation on the distribution and abundance of

- a chironomid, *Paraborniola tonnoiri* Freeman. Australian Journal of Zoology 22:71–89.
- Jutson J T 1934 The physiography (geomorphology) of Western Australia. Geological Survey of Western Australia, Perth. Bulletin 95.
- Lake P S, Bayly I A E & Morton D W 1989 The phenology of a temporary pond in western Victoria, Australia, with special reference to invertebrate succession. Archiv für Hydrobiologie 115:171–202.
- Lansbury I 1995 Notes on the Corixidae and Notonectidae (Hemiptera: Heteroptera) of southern Western Australia. Records of the Western Australian Museum 17:181–189.
- Maher M & Carpenter SM 1984 Benthic studies of waterfowl breeding habitat in south-western New South Wales. II Chironomid populations. Australian Journal of Marine and Freshwater Research 35:97–110.
- March F & Bass D 1995 Application of island biogeography theory to temporary pools. Journal of Freshwater Ecology 10:83–85.
- Petkovski T K 1973 Zur Cladoceran-fauna Australiens II Sididae und Macrothricidae. Acta Musei Macedonici Scientiarum Naturalium 13:161–192.
- Ranta E 1982. Animal communities in rock pools. Annales Zoologici Fennici 19:337–347.
- Sheldon A L 1984 Colonization dynamics of aquatic insects. In: The Ecology of Aquatic Insects (eds V H Resh & D M Rosenberg). Praeger, New York, 401–429.
- Smirnov N N & Bayly I A E 1995 New records and further description of *Macrothrix hardingi* Petkovski (Cladocera) from granite pools in Western Australia. Journal of the Royal Society of Western Australia 78:13–14.
- Smith L L 1941 Weather pits in granite of the southern Piedmont. Journal of Geomorphology 4:117–127.
- Timms B V 1981 Animal communities in three Victorian lakes of differing salinity. Hydrobiologia 81:181–193.
- Twidale C R & Corbin E M 1963 Gnammas. Revue de Géomorphologie dynamique 14:1–20.
- Wallwork J A 1981 A new aquatic oribatid mite from Western Australia (Acari: Cryptostigmata: Ameronothridae) Acarologia 22:333–339.
- Wiggins G B, Mackay R J & Smith I M 1980 Evolutionary and ecological strategies of animals in annual temporary pools. Archiv für Hydrobiologie, Supplement 58:97–206.
- Williams W D 1985 Biotic adaptations in temporary lentic waters, with special reference to those in semi-arid and arid regions. Hydrobiologia 125:85–110.
- Wolf E 1911 Phyllopoda. In: Die Fauna Südwest-Australiense Ergebnisse der Hamburger südwest-australischen Forschungsreise 1905 (eds W Michaelsen & R Hartmeyer). Vol. 3. G Fischer, Jena, 251–276.

Download Now. SaveSave Water harvesting from granite outcrops in Western For Later. 0 ratings0% found this document useful (0 votes). 25 views4 pages.Â Davis Gnammas, soaks and shallow wells generally have a (1977) reported that hundreds of shallow wells were limited water yield and are therefore seldom used for constructed as public water supplies in many of the public water supplies today. However, these sources of water were an important part of traditional Aboriginal culture. Supplies with larger water yields are more likely to Â© Royal Society of Western Australia 1997 involve the construction of walls, weirs or tanks. Granite Outcrops Symposium, 1996 Engineered water supplies consist of a storage reservoir. Granite outcrops of Western Australia are weathered landforms that occur throughout the state of Western Australia, composed primarily of the rock type granite. All recognised types of this landform can be observed, commonly as bornhardts, but also as inselbergs, castle koppies and Nubbins. Rising abruptly from the surrounding landscape they create a variety of microhabitats for plants, and provide seasonal resources and refuge for a range of animals. These areas thus have rich biodiversity and many Bayly IAE (1997) Invertebrates of temporary waters in gnammas on granite outcrops in Western Australia. J Royal Soc W Austral 80:167â€”172Google Scholar. Bayly IAE (2011) Australiaâ€™s granite wonderlands. Bas Publishing, MelbourneGoogle Scholar. Beisner BE, Hovius J et al (2006) Environmental productivity and biodiversity effects on invertebrate community invasibility.Â Morphometric heterogeneity of temporary pools in Malta and its effect on species richness: implications for management and restoration. Proc 6th EPCN Conference, Huesca, Spain, abstractGoogle Scholar. Buschke FT, Esterhuysen S et al (2012) The dynamics of mountain rock poolsâ€™are aquatic and terrestrial habitats alternative stable states? Acta Oecol 47:24â€”29CrossRefGoogle Scholar. Invertebrates of temporary waters in gnammas on granite outcrops in Western Australia. Article. Sep 1997.Â Nineteen pools distributed between nine different granite outcrops, all south of 32Â° s. in Western Australia, were studied from May to July 1977. Data on depth, conductivity, pH and temperature are presented. In all, 35 invertebrate taxa, mostly identified to species, were recorded. View. Show abstract. Habitat template for invertebrates on granite outcrops. Article. Jan 2000. J Roy Soc West Aust. Barbara York Main. This study considers the known occurrences of invertebrate taxa from granite outcrops of southern Western Australia. The microhabitats occupied are briefly described. A schematic h

Download now. SaveSave Water harvesting from granite outcrops in Western For Later. 0 ratings0% found this document useful (0 votes). 25 views4 pages.Â Davis Gnammas, soaks and shallow wells generally have a (1977) reported that hundreds of shallow wells were limited water yield and are therefore seldom used for constructed as public water supplies in many of the public water supplies today. However, these sources of water were an important part of traditional Aboriginal culture. Supplies with larger water yields are more likely to Â© Royal Society of Western Australia 1997 involve the construction of walls, weirs or tanks. Granite Outcrops Symposium, 1996 Engineered water supplies consist of a storage reservoir. Bayly, IAE (2002) The life of temporary waters in Australian gnammas (rockholes) Verh. Internat Verein Limnol 28:41â€“48Google Scholar. Beard JS (1990) Plant life of Western Australia.Â Withers PC, Edward DH (1998) Terrestrial fauna of granite outcrops in Western Australia. J R Soc West Aust 80:159â€“166Google Scholar. Yen AL (1987) A preliminary assessment of the correlation between plant, vertebrate and Coleoptera communities in the Victorian mallee. Bayly, I.A.E. (1997) Invertebrates of temporary waters in gnammas on granite outcrops in Western Australia. Journal of the Royal Society of Western Australia, 80, 167-172. Bayly, I.A.E. (1998) Beetle-mites that take to the water. Invertebrata, 12, 4. Berlese, A. (1886) Acari Myriopoda et Scorpiones Hucusque in Italia. Fasc. 33, no. 10. Pools of rainwater, known as gnammas,[1] often form on granite outcrops. These provide habitat to around 230 aquatic invertebrates, of which at least 50 are endemic to these pools,[2] and provide freshwater to other inhabitants and visitors. Gnamma on Mount Melville, Albany. The variable rock surfaces and soil types create niches for many plants. At least 1300 plant species occur on granite outcrops in Western Australia, many of which are endemic to these sites. YouTube Encyclopedic. 1/3. Invertebrate fauna and ecology of temporary pools on granite outcrops in southern Western Australia. Aust. J. mar. Freshwat. Res. 33: 599â€“606.CrossRefGoogle Scholar. Bayly, I. A. E. & P. Ellis, 1969.Â Seasonal fauna of some ephemeral saline waters in western Victoria with particular reference to *Parartemia zietziana*Sayce (Crustacea: Anostraca). Aust. J. mar. Freshwat. Res. 27: 1 â€“ 22.CrossRefGoogle Scholar. Geddes, M. C., 1983.